


ATROPINE SULFATE

Anticholinergic/(parasympatholitic) 0,5 mg

Anticholinergic/(parasympatholitic) 1 mg



10 syringes per box
Supplier number: FG00111AGT

7" / 18,5 cm


2,75" / 7,3 cm



10 syringes per box
Supplier number: FG00110AGT

7" / 18,5 cm

2,75" / 7,3 cm



AGUETTANT CANADA INC.

Tel.: 514 772-6294 | Toll-free: 1 833 772-6294

Fax: 1 888 772-0380

injectables@aguettant.ca | www.aguettant.ca

ATROPINE SULFATE


Anticholinergic/(parasympatholitic)

Ready to use

No preservative agent |

Shelf life of 3 years | Safe and sterile

Double gradation



Luer lock compatible

Latex free

Photos for illustration purposes only.

CONCENTRATIONS:

Atropine Sulfate (0.1 mg / mL for 0.5 mg / 5 mL)

Atropine Sulfate (0.2 mg / mL for 1 mg / 5 mL)

ROUTES OF ADMINISTRATIONS:

Intravenous, intramuscular and subcutaneous

DOSAGE FORMS:

Sterile solutions of Atropine (0.1 mg / mL et 0.2 mg / mL)

NON MEDICINAL INGREDIENTS:

Sodium Chloride (for tonicity), water for injection, Hydrochloride Acid (for pH adjustment)

STORAGE:

Store syringe in original blister between 15-30°C. Avoid excessive heat. Protect from freezing.


HANDLING INSTRUCTION:

The unopened and undamaged blister package is sterile and should not be opened before use. The solution should be visually inspected for absence of particles and discoloration prior to administration. Do not use if the mixture shows haziness or if the tamper evident seal on the syringe is broken. Use immediately after opening. Discard unused portion.

	CPDN #	McKesson #
ATROPINE 0,5 mg	901100	172345
ATROPINE 1 mg	901117	171425
EPINEPHRINE 1 mg	902696	172711
LIDOCAINE 1% 10 mL	902429	179860

	CPDN #	McKesson #
LIDOCAINE 2% 5 mL	902375	179859
LIDOCAINE 2% 10 mL	902443	179832
PHENYLEPHRINE 500 mcg	901186	103187


Metabisulfite-free/
Sulfite-free




1. Withdraw the pre-filled syringe from the sterile blister.




2. The sterilization process may have caused the plunger to stick to the pre-filled syringe. **Press** the plunger to release it from the wall.




3. Twist off the end cap to break the tamper evident seal. In order to avoid contamination, do not touch the exposed luer connection.



4. Check the pre-filled syringe seal tip has been completely removed. If not, replace the cap and twist again.



5. Expel the air by gently pushing the plunger.



6. Connect the pre-filled syringe to access device or the needle. Push the plunger slowly to inject the required volume.



PRODUCT LIST

Solutions for injection in pre-filled syringes

Products	Dosing/ Volume	DIN	Shelf life
Atropine	0.5 mg / 5 mL	02426269	36 months
Atropine	1 mg / 5 mL	02426277	36 months
Epinephrine	1 mg / 10 mL	02521148	24 months
Lidocaine 1%	100 mg / 10 mL	02500647	36 months
Lidocaine 2%	100 mg / 5 mL	02522268	36 months
Lidocaine 2%	200 mg / 10 mL	02500655	36 months
Phenylephrine	500 mcg / 10 mL	02454599	36 months