

PRESCRIBING INFORMATION

EPINEPHRINE INJECTION

0.1 mg / mL (Single-Use Syringes)

Sterile Solution
House standard

(Sympathomimetic)

Laboratoire Aguetant
1 rue Alexander Fleming
69007 Lyon
France

Date of preparation: 07-10-2021

Imported to Canada by:
Aguettant Canada Inc.
1470 Peel Suite A-152, Montreal QC, H3A 1T1

Control No.: 235107

NAME OF DRUG

EPINEPHRINE INJECTION

House Standard

0.1 mg / mL (Single-Use Syringes)

Sterile Solution

THERAPEUTIC CLASSIFICATION

Sympathomimetic

Epinephrine Injection is a parenteral adrenergic (sympathomimetic) agent and cardiac stimulant.

ACTION AND CLINICAL PHARMACOLOGY

The actions of epinephrine resemble the effects of stimulation of adrenergic nerves. It acts on both alpha and beta receptor sites of sympathetic effector cells. Its most prominent actions are on the beta receptors of the heart, vascular and other smooth muscle. When given by rapid intravenous injection, it produces a rapid rise in blood pressure, mainly systolic, by (1) direct stimulation of cardiac muscle which increases the strength of ventricular contraction, (2) increasing the heart rate and (3) constriction of the arterioles in the skin, mucosa and splanchnic areas of the circulation.

When given by slow intravenous injection, epinephrine usually produces only a moderate rise in systolic and a fall in diastolic pressure. Although some increase in pulse pressure occurs, there is usually no great elevation in mean blood pressure. Accordingly, the compensatory reflex mechanisms that come into play with a pronounced increase in blood pressure do not antagonize the direct cardiac actions of epinephrine as much as with catecholamines that have a predominant action on alpha receptors.

Total peripheral resistance decreases by action of epinephrine on beta receptors of the skeletal muscle vasculature and blood flow is thereby enhanced. Usually this vasodilator effect of the drug on the circulation predominates so that the modest rise in systolic pressure which follows slow injection or absorption is mainly the result of direct cardiac stimulation and increase in cardiac output. In some instances, peripheral resistance is not altered or may even rise owing to a greater ratio of alpha to beta activity in different vascular areas.

Epinephrine relaxes the smooth muscles of the bronchi and iris and is a physiologic antagonist of histamine. The drug also produces an increase in blood sugar and glycogenolysis in the liver.

Pharmacokinetics

Intravenous injection produces an immediate and intensified response. Following intravenous injection, epinephrine disappears rapidly from the blood stream.

Epinephrine is rapidly inactivated in the body and is degraded by enzymes in the liver and other tissues. The larger portion of injected doses is excreted in the urine as inactivated compounds and the remainder either partly unchanged or conjugated.

The drug becomes fixed in the tissues and is inactivated chiefly by enzymatic transformation to metanephrine or normetanephrine, either of which is subsequently conjugated and excreted in the urine in the form of sulfates and glucuronides. Either sequence results in the formation of 3-methoxy-4-hydroxy-mandelic acid (vanillyl-mandelic acid; VMA) which also is detectable in the urine.

Sodium chloride added to render the solution isotonic for injection of the active ingredient is present in amounts insufficient to affect serum electrolyte balance of sodium (Na^+) and chloride (Cl^-) ions.

INDICATIONS AND CLINICAL USE

In general, the most common uses of parenteral epinephrine are to relieve respiratory distress due to bronchospasm, to provide rapid relief of hypersensitivity (anaphylactic or anaphylactoid) reactions to drugs, animal serums and other allergens. Its cardiac effects may be of use in restoring cardiac rhythm in cardiac arrest due to various causes, and attacks of transitory atrioventricular (A-V) heart block and syncopal seizures (Stokes-Adams syndrome), but it is not used in cardiac failure or in hemorrhagic, traumatic, or cardiogenic shock.

Epinephrine is used as a hemostatic agent.

It is also used in treating mucosal congestion of hay fever, rhinitis, and acute sinusitis; to relieve bronchial asthmatic paroxysms; in syncope due to complete heart block or carotid sinus hypersensitivity; for symptomatic relief of serum sickness, urticaria, angioneurotic edema; for resuscitation in cardiac arrest following anesthetic accidents; in simple (open angle) glaucoma; for relaxation of uterine musculature and to inhibit uterine contractions. (See **CONTRAINDICATIONS.**)

CONTRAINDICATIONS

Epinephrine is contraindicated in patients with known hypersensitivity to sympathomimetic amines, in patients with angle closure glaucoma, and patients in shock (nonanaphylactic). It should not be used in patients anesthetized with agents such as cyclopropane or halothane as these may sensitize the heart to arrhythmic action of sympathomimetic drugs.

Epinephrine should not ordinarily be used in cases where vasopressor drugs may be contraindicated, e.g. in thyrotoxicosis, diabetes, patients receiving MAO inhibitors, in obstetrics when maternal blood pressure is in excess of 130/80 or during labour, and in hypertension and other cardiovascular disorders.

WARNINGS

Inadvertently induced high arterial blood pressure may result in angina pectoris, aortic rupture or cerebral hemorrhage.

Epinephrine may induce potentially serious cardiac arrhythmias in patients not suffering from heart disease and patients with organic heart disease or who are receiving drugs that sensitize the myocardium.

Parenterally administered epinephrine initially may produce constriction of renal blood vessels and decrease urine formation, and large doses may cause complete renal shutdown.

Epinephrine and Beta-blockers

There may be increased difficulty in treating an allergic-type reaction in patients on beta-blockers. In these patients, the reaction may be more severe due to pharmacologic effects of the beta-blockers and problems with fluid changes (see **PRECAUTIONS-Drug Interactions**).

PRECAUTIONS

Although epinephrine can produce ventricular fibrillation, its actions in restoring electrical activity in asystole and in enhancing defibrillation of the fibrillating ventricle are well documented. The drug, however, should be used with caution in patients with ventricular fibrillation.

In patients with prefibrillatory rhythm, intravenous epinephrine must be used judiciously, with extreme caution, because of its excitatory action on the heart. Since the myocardium is sensitized to this action of the drug by many anesthetic agents, epinephrine may convert asystole to ventricular fibrillation if used in the treatment of anesthetic cardiac accidents.

Epinephrine should be used cautiously in the elderly and in patients with hyperthyroidism, hypertension, diabetes and cardiac diseases/arrhythmias. Patients with long-standing bronchial asthma and emphysema, who have developed degenerative heart disease, should be administered the drug with extreme caution.

Fatalities may also result from pulmonary edema because of the peripheral constriction and cardiac stimulation produced. Rapidly acting vasodilators such as nitrites, or alpha-blocking agents may counteract the marked pressor effects of epinephrine.

Drug Interactions

Beta-blockers

There may be increased difficulty in treating an allergic-type reaction in patients on beta-blockers. In these patients, the reaction may be more severe due to pharmacologic effects of the beta-blockers and problems with fluid changes. Epinephrine should be administered with caution, since it may not have its usual effects in the treatment of anaphylaxis. On the one hand, larger doses of epinephrine may be needed to overcome the bronchospasm, while on the other, these doses can be associated with excessive alpha-adrenergic stimulation with consequent hypertension, reflex bradycardia and heart-block and possible potentiation of bronchospasm. Alternatives to the use of large doses of epinephrine include vigorous supportive care such as fluids and the use of beta agonists including parenteral salbutamol or isoproterenol to overcome bronchospasm and norepinephrine to overcome hypotension.

Sympathomimetic Drugs

Epinephrine should not be administered concomitantly with other sympathomimetic drugs (such as isoproterenol) because of possible additive effects and increased toxicity. Combined effects may induce serious cardiac arrhythmias. They may be administered alternately when the preceding effect of another such drug has subsided.

Cyclopropane or Halogenated Hydrocarbons

Administration of epinephrine to patients receiving cyclopropane or halogenated hydrocarbon general anesthetics such as halothane, which sensitize the myocardium, may induce cardiac arrhythmias (see **CONTRAINDICATIONS**). When encountered, such arrhythmias may respond to administration of a beta-adrenergic blocking drug.

Diuretics

Diuretic agents may decrease vascular response to pressor drugs such as epinephrine.

Guanethidine

Epinephrine may antagonize the neuron blockade produced by guanethidine, resulting in decreased antihypertensive effect and requiring increased dosage of the latter.

MAO Inhibitors

All vasopressors should be used cautiously in patients taking monoamine oxidase (MAO) inhibitors (see **CONTRAINDICATIONS**).

Others

The effects of epinephrine may be potentiated by tricyclic antidepressants; certain antihistamines, e.g. diphenhydramine, tripeleminamine, chlorpheniramine; and sodium levothyroxine.

Epinephrine also should be used cautiously with other drugs (e.g. digitalis glycosides) that sensitize the myocardium to the actions of sympathomimetic drugs.

Pregnancy

Epinephrine has been shown to be teratogenic in rats when given in doses about 25 times the human doses. It is not known whether epinephrine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Epinephrine should be given to a pregnant woman only if clearly needed.

Labor and Delivery

Parenteral administration of epinephrine, if used to support blood pressure during low or other spinal anesthesia for delivery, can cause acceleration of fetal heart rate and should not be used in obstetrics when maternal blood pressure exceeds 130/80 (see **CONTRAINDICATIONS**). Epinephrine may delay the second stage of labour.

ADVERSE REACTIONS

Transient and minor side effects of anxiety, headache, fear and palpitations may occur with systemic therapeutic doses, especially in hyperthyroid individuals. Adverse effects, such as cardiac arrhythmias and excessive rise in blood pressure, may also occur with systemic therapeutic doses or inadvertent overdosage. Other adverse reactions include: cerebral hemorrhage, hemiplegia, subarachnoid hemorrhage, anginal pain in patients with angina pectoris, anxiety, restlessness, throbbing headache, tremor, weakness, dizziness, pallor and respiratory difficulty.

SYMPTOMS AND TREATMENT OF OVERDOSAGE

Erroneous administration of large doses of epinephrine may lead to precordial distress, vomiting, headache, dyspnea, as well as unusually elevated blood pressure (see **WARNINGS**). Toxic effects of overdosage can be counteracted by injection of an alpha-adrenergic blocker and a beta-adrenergic blocker. In the event of a sharp rise in blood pressure, rapid acting vasodilators such as the nitrites, or alpha-adrenergic blocking agents can be given to counteract the marked pressor effect of large doses of epinephrine.

For management of a suspected drug overdose, contact your regional poison control centre.

DOSAGE AND ADMINISTRATION

Epinephrine Injection is administered by the following routes: intravenous, and via endotracheal tube in the bronchial tree.

Hypersensitivity reaction

Adults:

For intramuscular administration, a 1 mg/ml solution should be used. Epinephrine Injection 1 mg/10 ml is not recommended for intramuscular or subcutaneous use in acute anaphylaxis

In severe anaphylactic shock, intravenous administration may be necessary since absorption of the drug may be impaired with subcutaneous or intramuscular administration. If necessary, 0.1 to 0.25 mg of epinephrine [1 to 2.5 mL of a 0.1 mg/mL solution] may be administered intravenously slowly (over 5 to 10 minutes) and repeated every 5 to 15 minutes as necessary. Start with small doses and increase if required.

Cardiac Resuscitation

Adults:

A dose of 0.5 mg intravenous [range 0.1 to 1 mg, usually as 1 to 10 mL of a 0.1 mg/mL solution]. Intravenous doses may be repeated every 5 minutes if needed.

Children:

The usual **pediatric** intravenous dose is 0.01 mg/kg [0.1 mL/kg of 0.1 mg/mL solution]. Intravenous doses may be repeated every 5 minutes if needed. Epinephrine Injection 1 mg/10 ml is not appropriate to deliver a dose of less than 0.5 ml. This product is not suitable for children less than 5 kg.

The usual **neonatal** intravenous dose is 0.01 to 0.03 mg/kg [0.1 to 0.3 mL/kg of a 0.1 mg/mL solution]. Intravenous doses may be repeated every 5 minutes if necessary. Epinephrine Injection 1 mg/10 ml is not appropriate to deliver a dose of less than 0.5 ml. This product is not suitable for neonates less than 2 kg.

Endotracheal Dosage

Alternatively, as a means for advanced cardiac life support, when vascular access is hampered and patients intubated, epinephrine can be administered via the endotracheal tube directly into the bronchial tree.

Adults: 1 mg [10 mL of a 0.1 mg/mL solution].

Children: 0.01 mg/kg [0.1 mL/kg of a 0.1 mg/mL solution]. Epinephrine Injection 1 mg/10 ml is not appropriate to deliver a dose of less than 0.5 ml. This product should NOT be used in children less than 5 kg.

Neonates: 0.01 to 0.03 mg/kg [0.1 to 0.3 mL/kg of a 0.1 mg/mL solution]. Epinephrine Injection 1 mg/10 ml is not appropriate to deliver a dose of less than 0.5 ml. This product should NOT be used in neonates less than 2 kg.

AVAILABILITY OF DOSAGE FORMS

Epinephrine Injection is a sterile, nonpyrogenic solution of epinephrine in water for injection.

Each milliliter of the 0.1 mg / mL solution contains epinephrine 0.1 mg (as epinephrine bitartrate) in water for injection, sodium chloride 9 mg. May contain additional sodium hydroxide and/or hydrochloric acid for pH adjustment - pH (3.0 to 3.4).

The solution contains no bacteriostatic or antimicrobial agent and is intended for use as a single-dose injection. When smaller doses are required, the unused portion should be discarded.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Storage: Store between 15 and 25°C. Store in the overwrap provided. Protect from light, oxygen, freezing and excessive heat.

RETAIN PRODUCT IN CARTON UNTIL READY FOR USE.

NOTE: DO NOT USE THE INJECTION IF IT IS COLOURED OR IF IT CONTAINS A PRECIPITATE. DO NOT ADMINISTER UNLESS SOLUTION IS CLEAR AND SEAL IS INTACT. DISCARD UNUSED PORTION.

Container	Size (mL)	Concentration (mg / mL)	Total content (mg Epinephrine)	Needle
Pre-filled syringe (individually packaged in a blister overwrapped in an aluminium bag with an oxygen absorber)	10	0.1 (Adult or Pediatric)	1.00	N.A.

REFERENCES

1. Pfizer Canada Inc., Prescribing Information for EPINEPHRINE INJECTION USP (1 mg/mL Ampoules and 0.1 mg/mL Single-Use Syringes), Control No.: 213923, Date of Revision: March 18, 2021.